




2010

BARLEY QUALITY

REPORT

2010 IDAHO BARLEY QUALITY REPORT



Summary

Idaho barley production totaled 43.24 million bushels from 470,000 harvested acres in 2010, representing a decrease of 10% in output from the previous year. Quality was mostly good overall.


Approximately 65% of Idaho's barley sampled received a USDA/FGIS grade of #1. There were 506 barley samples collected from Idaho elevators, commodity brokers and farmers. The samples were separated by class, variety and region. They were graded at a licensed Federal Grain Inspection Service (FGIS) facility.

Average weight was slightly lower than last year at 49.6 pounds per bushel (lb/bu). Eighty percent of the 2-row malting barley crop tested 48 lb/bu or better and seventy-seven percent of the 6-row malting barley crop tested 48 lb/bu or better. Overall thins averaged 2.09% and plumps averaged 92.4%. Protein averaged 11.3% statewide and moisture averaged 10.5%. The average for skinned and broken kernels was 1.5%.

Average Statewide Barley Quality

Type	# Samples	Test Wt. ¹		1000 Knl (grams)	Moist %	Thin ² %	Protein	Sound ³	Skn/Br ⁴ %	% Plump	Average
		(lb/bu)	(kg/hl)								
2-R Feed	72	48.43	62.38	41.84	10.41	3.13	11.93	99.27	1.35	91.37	1.57
6-R Feed	9	49.10	63.24	42.60	9.99	3.04	11.34	99.06		97.20	1.22
2-R Malting	103	50.66	65.25	42.47	10.49	3.46	10.84	99.59	1.58	92.69	1.38
6-R Malting	322	49.91	64.28	36.29	10.78		11.84	99.70	1.74	97.08	1.24
2010 Avg.	506	49.59	63.88	41.35	10.48	2.09	11.31	99.96	1.47	92.44	1.43
2009 Avg.	843	49.76	64.09	41.10	10.76	1.98	11.81	99.52	1.90	83.14	1.68
5 Yr Avg.	425	49.35	63.59	39.26	9.72	3.63	11.59	99.46	1.56	86.22	1.51

2010 IDAHO BARLEY VARIETIES MALTING VS FEED



Top Ten Idaho Varieties

Variety	Planted Acres	Hectares	% of Prod.
Conrad (2-Row Malt)	102,700	41,562	23.7%
AC Metcalfe (2-Row Malt)	85,000	34,399	19.6%
Moravian 69 (2-Row Malt)	54,500	22,056	12.6%
Baronesse (2-Row Feed)	49,100	19,870	11.3%
Merit (2-Row Malt)	33,000	13,355	7.6%
Harrington (2-Row Malt)	32,200	13,031	7.4%
Legacy (6-Row Malt)	20,600	8,337	4.8%
Copeland (2-Row Malt)	16,500	6,677	3.8%
Criton (2-Row Feed)	14,300	5,787	3.3%
Charles (2-Row Malt)	11,300	4,573	2.6%
Other Malt	2,600	1,052	0.6%
Other Feed	11,700	4,735	2.7%
TOTAL:	433,500	175,435	100%


Source: 2010 IBC/Idaho Ag Statistics Service

Source: 2010 IBC/Idaho Ag Statistics Service

2010 Idaho Barley Quality By Variety


Variety	Type/Use	# Samples	Test Wt. (lbs/bu)	1000 Knl (grams)	Moisture %	Thin %	Protein %	Sound %	Skn/Br%	Plump%	Average
Legacy	2-Row Malt	109	49.83	36.95	10.93	1.51	11.65	99.73	2.22	97.37	1.19
Conrad	2-Row Malt	96	50.69	42.66	10.69	1.79	11.89	99.58	1.11	94.58	1.13
AC Metcalfe	2-Row Malt	73	49.72	45.93	10.92	1.55	12.19	99.73	1.50	90.23	1.88
Harrington	2-Row Malt	62	49.65	40.50	10.61	1.94	11.69	99.54	1.78	88.92	2.08
Moravian 69	2-Row Malt	33	51.09	45.12	9.78	1.98	10.66	99.81	1.32	90.68	1.00
Copeland	2-Row Malt	18	45.33	30.50	10.53	2.29	10.24	88.83	1.35	88.35	1.91
Tradition	6-Row Malt	16	49.53	33.30	10.87	1.56	11.98	99.60	2.34	89.49	1.33
Merit	2-Row Malt	14	49.14	46.02	9.95		10.53	99.75	1.49	89.73	1.46

Idaho Barley Test Weight Distribution 2009-2010



2010 Regional Quality

North Idaho



Barley Production

	<u>2009</u>	<u>2010</u>
<u>Harvested (Acres)</u>	40,800	32,900
(Hectares)	16,483	13,314
% Idaho Acreage	8%	7%


Source: Idaho Ag. Statistical Service/IBC

Barley Varieties Planted

Champion	26.9%
Copeland	19.7%
Baronesse	10.3%
Camas	5.4%
Conrad	4.9%
Steptoe	2.0%
Other Malting	0.6%
Other Feed	10.0%

Type	# Samples	Test Wt. ¹ (lb/bu) (kg/hl)	1000 Knl (grams)	Moist %	Thin ² %	Protein	Sound ³	Skn/Br ⁴ %	% Plump ⁵	Average
2-R Feed	8	49.27	63.46	10.53		10.99	99.69			2.00
6-R Feed	6	49.61	63.89	40.36	10.63	11.62	99.80	1.80	88.90	1.83
2010 Avg.	14	49.41	63.64	40.36	10.58	11.26	99.74	1.80	88.90	1.93
2009 Avg.	14	50.86	65.50	40.30	10.39	12.00	99.53	2.15	90.79	1.14
5 Yr Avg.	15	48.13	59.65	37.52	8.86	11.39	91.16	1.77	76.46	1.31

Southwest Idaho



Barley Production

	<u>2009</u>	<u>2010</u>
<u>Harvested (Acres)</u>	8,160	5,640
(Hectares)	3,304	2,282
% Idaho Acreage	1.6%	1.2%

Source: Idaho Ag. Statistical Service/IBC

Barley Varieties Planted

Steptoe	28.3%
Baronesse	15.0%
Idagold	11.7%
Millennium	5.0%
Other Feed	30.0%


Type	# Samples	Test Wt. ¹ (lb/bu) (kg/hl)	1000 Knl (grams)	Moist %	Thin ² %	Protein	Sound ³	Skn/Br ⁴ %	% Plump ⁵	Average
2-R Feed	6	51.63	66.50	0.00	11.52	10.67	99.55			1.00
6-R Feed	2	49.45	63.69	0.00	10.45	10.20	99.20			1.00
2010 Avg.	8	51.09	65.80	0.00	11.25	10.55	99.46			1.00
2009 Avg.	8	47.22	60.82	0.00	9.28	11.51	99.28			1.53
5 Yr Avg.	9	44.70	53.90	0.00	7.53	9.71	83.85			1.28

¹ Weighted average by regional production

² % Thins: 2-rowed kernels passing through a 5/64 x 3/4 in. (2.2 x 19 mm) slotted sieve
6-rowed kernels passing through a 5/64 x 3/4 in. (2.0 x 19 mm) slotted sieve

2010 Regional Quality

South Central Idaho



Barley Production

	2009	2010
Harvested (Acres)	127,500	125,020
(Hectares)	51,619	50,594
% Idaho Acreage	25%	26.6%


Source: Idaho Ag. Statistical Service/IBC

Barley Varieties Planted

Moravian 69	34.5%
Conrad	16.0%
Criton	10.0%
Charles	8.4%
Baronesse	4.9%
Idagold	4.2%
AC Metcalfe	2.6%
Other Malting	1.5%
Other Feed	2.1%

Type	# Samples	Test Wt. ¹		1000 Knl (grams)	Moist %	Thin ² %	Protein	Sound ³	Skn/Br ⁴ %	% Plump ⁵	Average
		(lb/bu)	(kg/hl)								
2-R Feed	22	50.67	65.27		9.33	2.96	13.58	99.81			1.27
6-R Feed	4	49.65	63.95		9.28	4.83	10.53	99.80			1.25
2-R Malting	18	52.44	67.55	43.21	9.82	2.34	10.98	99.56	1.41	91.50	2.20
6-R Malting	1	48.20	62.08	34.60	9.50	1.50	9.60	99.70	1.70	84.00	1.00
2010 Avg.	27	51.15	65.88	43.21	9.91	3.01	11.20	99.75	1.41	91.50	1.48
2009 Avg.	45	50.05	64.47	38.79	9.40	2.91	11.17	99.79	1.69	88.04	1.20
5 Yr Avg.	67.6	54.93	70.75	40.73	10.22	3.51	10.67	97.66	1.42	90.24	1.34

Southeast Idaho



Barley Production

	2009	2010
Harvested (Acres)	333,540	306,440
(Hectares)	135,036	124,012
% Idaho Acreage	65.4%	65.2%

Source: Idaho Ag. Statistical Service/IBC

Barley Varieties Planted


Conrad	24.4%
AC Metcalfe	24.3%
Baronesse	11.7%
Merit	9.6%
Harrington	8.5%
Legacy	6.0%
Copeland	2.9%
Tradition	2.6%
Moravian 69	2.3%
Other Malting	0.1%
Other Feed	1.1%

Type	# Samples	Test Wt. ¹		1000 Knl (grams)	Moist %	Thin ² %	Protein	Sound ³	Skn/Br ⁴ %	% Plump ⁵	Average
		(lb/bu)	(kg/hl)								
2-R Feed	52	47.58	61.29	41.38	10.18	2.95	12.21	99.29	1.27	87.15	1.63
6-R Feed	3	48.17	62.04	42.60	11.23	3.33	13.03	97.97	0.90	97.20	1.33
2-R Malting	70	50.06	64.48	42.30	10.77	3.63	10.88	98.03	1.65	93.63	1.42
6-R Malting	322	49.91	64.28	36.29	10.78	1.45	11.84	99.70	1.80	92.86	1.24
2010 Avg.	447	48.93	63.02	40.64	70.87	2.84	11.99	98.75	1.40	92.71	1.41
2009 Avg.	776	48.56	62.54	38.01	10.78	2.81	11.78	99.37	1.47	85.92	1.49
5 Yr Avg.	326	48.62	62.62	55.52	9.87	3.28	11.96	98.61	1.56	85.17	1.55

³ % Sound = 100 % dockage free barley (minus) % damaged kernels, % foreign material, % other grains, % wild oats



⁴ % Skinned and broken kernels ⁵ % Plumps: Kernels retained on or above 6/64 x 3/4 in. (2.4 x 19 mm) slotted sieve

Idaho Barley Protein Distribution 2009-2010



2-Row Malting Barley Protein

6-Row Malting Barley Protein



METHODS EMPLOYED AND DEFINITION OF TERMS AND SYMBOLS

TEST WEIGHT - Test weight was determined on dockage free barley using a Seedbuero Model #8850 computer grain scale. Test weight was expressed in pounds per bushel. In the event sample moisture exceeded 13.5%, the sample was allowed to air-dry prior to test weight determination.

MOISTURE - Moisture expressed as % of total weight was determined by using a Dicky-john gac2100.

PROTEIN - Percent total protein calculated on a dry basis was determined by infratech model 1241 whole grain analyzer using ANN calibration.

PLUMP BARLEY - Barley that remains on top of a 6/64 x 3/4 slotted-hole sieve after sieving.

THIN BARLEY - For the class Barley¹ and 6-row barley which passes through a 5/64 x 3/4 slotted-hole sieve and 2-row malting barley² which passes through a 5.5/64 x 3/4 slotted-hole sieve after sieving.

¹ The class Barley is a mixture of 10% or more of 2-row barley or 6-row barley.

² 2-row malting barley is sieved on a 5.5/64 x 3/4 slotted-hole sieve which would be higher thins.

USDA/FGIS

United States Grades and Grade Requirements

§ 810.204 Grades and grade requirements for the subclasses Six-rowed Malting Barley and Six-rowed Blue Malting Barley.

Grade	Minimum limits of -			Maximum Limits of -				
	Test Weight Per/bu.	Suitable Malting Types	Sound Barley ¹	Damaged Kernels ¹	Foreign Material	Other Grains	Skinned & Broken Kernels	Thin Barley ²
	<i>Pounds</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
U.S. No. 1	47.0	95.0	97.0	2.0	.5	2.0	4.0	7.0
U.S. No. 2	45.0	95.0	94.0	3.0	1.0	3.0	6.0	10.0
U.S. No. 3	43.0	95.0	90.0	4.0	2.0	5.0	8.0	15.0
U.S. No. 4	43.00	95.0	87.0	5.0	3.0	5.0	10.0	15.0

¹ Injured-by-frost kernels and injured-by-mold kernels are not considered damaged kernels or considered against sound barley..

² Use the 5/64 x 3/4 slotted-hole sieve

Note: Malting barley shall not be infested in accordance with § 810.07(b) and shall not contain any special grades as defined in § 810.206. Six-rowed Malting barley and Six-rowed Blue Malting barley varieties not meeting the requirements of this section shall be graded in accordance with standards established for the class Barley.

§ 810.205 Grades and grade requirements for Two-rowed Malting Barley.

Grade	Minimum limits of -			Maximum Limits of -			
	Test Weight Per/bu.	Suitable Malting Types	Sound Barley ¹	Wild Oats	Foreign Material	Skinned & Broken Kernels	Thin Barley ²
	<i>Pounds</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
U.S. No. 1	50.0	97.0	98.0	1.0	0.5	5.0	5.0
U.S. No. 2	48.0	97.0	98.0	1.0	1.0	7.0	7.0
U.S. No. 3	48.0	95.0	96.0	2.0	2.0	10.0	10.0
U.S. No. 4	48.0	95.0	93.0	3.0	3.0	10.0	10.0

¹ Injured-by-frost kernels and injured-by mold kernels are not considered damaged kernels or considered against sound barley

² Use the 5.5/64 x 3/4 slotted-hole sieve.

Note: Malting barley shall not be infested in accordance with § 810.07(b) and shall not contain any special grades as defined in § 810.206. Two-rowed Malting barley varieties not meeting the requirements of this section shall be graded in accordance with standards established for the class Barley.

§ 810.206 Grades and grade requirements for Barley.

Grade	Minimum limits of -			Maximum Limits of -			
	Test Weight Per/bu.	Sound Barley ¹	Damaged Kernels ¹	Heat Damaged Kernels	Foreign Material	Skinned & Broken Kernels	Thin Barley ²
	<i>Pounds</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>	<i>Percent</i>
U.S. No. 1	47.0	97.0	2.0	0.2	1.0	4.0	10.0
U.S. No. 2	45.0	94.0	4.0	0.3	2.0	8.0	15.0
U.S. No. 3	43.0	90.0	6.0	0.5	3.0	12.0	25.0
U.S. No. 4	40.0	85.0	8.0	1.0	4.0	18.0	35.0
U.S. No. 5	36.0	75.0	10.0	3.0	5.0	28.0	75.0

U.S. Sample grade shall be barley that:

(a) Does not meet the requirements for the grades 1, 2, 3, 4, or 5: or

(b) Contains 8 or more stones or any number of stones which have an aggregate weight in excess of 0.2 percent of the sample weight, 2 or more pieces of glass, 3 or more crotalaria seeds (*Crotalaria* spp.), 2 or more castor beans (*Ricinus communis* L), 4 or more particles of unknown foreign substance(s) or commonly recognized harmful or toxic substance(s), 8 or more cocklebur (*Xanthium* spp.) or similar seeds singly or in combination, 10 or more rodent pellets, bird droppings, or equivalent quantity of other animal filth per 1-1/8 to 1-1/4 quarts of barley: or (c) Has a musty, sour, or commercially objectionable foreign odor (except smut or garlic odor); or (d) Is heating or otherwise of distinctly low quality.

¹ Includes heat damaged kernels. Injured-by-frost kernels and injured-by-mold kernels are not considered damaged kernels or considered against sound barley..

² Use the 5/64 x 3/4 slotted-hole sieve

METRIC CONVERSION TABLES FOR BARLEY

U.S. to Metric	
1 Wet Quart (Qt)	= 0.9464 Liter (L)
1 Dry Quart	= 1.10 Liters
1 Wet Gallon (Gal)	= 3.785 Liters
1 Wet Gallon	= 0.0378 Hectoliter (Hl)
1 Dry Gallon	= 4.404 Liters
1 Dry Gallon	= 0.044 Hectoliter
1 Bushel	= 0.352 Hectoliter
1 Acre (A)	= 0.404694 Hectare (Ha)
1 Pound (Lb)	= 0.4536 Kilogram (Kg)
1 Mile (Mi)	= 1.610 Kilometers (Km)
1 Bu. Barley	= .021772 Metric Ton
1 Beer Barrel	= 31.0 Gallons
Lb/Bu x 1.288	= Kg/Hl
Bu/A x 0.869	= Hl/Ha
Bu/A x .05380	= MT/Ha

Metric to U.S.	
1 Liter	= 1.0567 Wet Quarts
1 Liter	= 0.098 Dry Quart
1 Liter	= .2642 Wet Gallon
1 Hectoliter	= 26.42 Wet Gallons
1 Liter	= .227 Dry Gallon
1 Hectoliter	= 22.70 Dry Gallons
1 Hectoliter	= 2.838 Bushels
1 Hectare	= 2.471 Acres
1 Kilogram	= 2.2046 Pounds
1 Kilometer	= 0.62 Mile
1 Metric Ton	= 45.9296 Bu. of Barley
1.1734 Hectoliters	= 1 Beer Barrel
Kg/Hl x .7764	= Lb/Bu
Hl/Ha x 1.15	= Bu/A
MT/Ha x 18.59	= Bu/A

STANDARDS

	Lbs/Bu	Kg/Hl
Malt	34	43.79
Barley	48	61.82

ACKNOWLEDGMENTS

The Idaho Barley Commission would like to thank the barley producers, elevator operators and commodity dealers who participated in our barley quality survey. Their combined efforts were responsible for the success of this survey.

Additional copies of this report are available by contacting the Idaho Barley Commission.



821 W State Street
 Boise, ID 83702
 Ph: (208) 334-2090
 Fax: (208) 334-2335
kolson@idahobarley.org